

WEST Search History

DATE: Monday, May 08, 2006

Hide?	Set Name	Query	Hit Count
		<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI,TDBD; PLUR=YES; OP=AND</i>	
<input type="checkbox"/>	L4	L3 and extracellular	40
<input type="checkbox"/>	L3	(gdf-8 or myostatin) same activin receptor	78
<input type="checkbox"/>	L2	l1 and muscle	84
<input type="checkbox"/>	L1	activin receptor gdf-8	98

END OF SEARCH HISTORY



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Reg]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bc

Search PubMed for

Limits Preview/Index **History** Clipboard Details

About Entrez
NCBI Toolbar

Text Version

- Search History will be lost after eight hours of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.
- Click on query # to add to strategy

Entrez PubMed
Overview
Help | FAQ
Tutorials
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Special Queries
LinkOut
My NCBI

Related Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Search	Most Recent Queries	Time	Result
#16	Search activin type II receptor extracellular	11:44:48	105
#8	Search activin receptor fusion	11:25:11	80
#9	Search activin receptor fusion extracellular	11:17:24	13
#1	Search activin receptor and gdf-8	11:12:54	8
#4	Search activin receptor and muscle	11:12:21	82
#2	Search activin and gdf-8	11:04:06	18

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Apr 24 2006 06:33:44



Search for

Limits Preview/Index History Clipboard Details

Display Show Sort by Send to

All: 1 Review: 0

About Entrez
NCBI Toolbar

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorials
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Special Queries
LinkOut
My NCBI

Related Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

☐ 1: J Protein Chem. 1999 May;18(4):437-46.

[Related Articles, Links](#)



The disulfide bond arrangement in the extracellular domain of the activin type II receptor.

Fischer WH, Greenwald J, Park M, Craig AG, Choe S, Vale W.

The Clayton Foundation Laboratories for Peptide Biology, The Salk Institute, La Jolla, California 92037, USA. Fischer@salk.edu

The initial step in the signaling cascade of the growth factor activin involves its binding to the extracellular domain of the activin type II receptor. This receptor domain contains 10 cysteine residues which are engaged in intramolecular disulfide bonds. To elucidate the structural framework of this domain we have characterized its disulfide-bonding pattern using an extracellular fragment of the receptor which binds activin A with high affinity. By combining proteolysis with mass spectroscopy and chemical sequence analysis, the disulfide connectivity was determined to be as follows: C1-C3, C2-C4, C5-C8, C6-C7, and C9-C10. A similar disulfide arrangement occurs in a family of snake toxins for which the three-dimensional structure is known.

PMID: 10449041 [PubMed - indexed for MEDLINE]

Display Show Sort by Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Apr 24 2006 06:33:44



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Register]

[All Databases](#)[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[OMIM](#)[PMC](#)[Journals](#)[Books](#)

Search for

[Limits](#)[Preview/Index](#)[History](#)[Clipboard](#)[Details](#)

Display Show Sort by Send to

All: 1 Review: 0

About Entrez
NCBI Toolbar

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorials
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Special Queries
LinkOut
My NCBI

Related Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

☐ 1: [Biochem Biophys Res Commun. 1992 Apr 15;184\(1\):310-6.](#)

[Related Articles](#),
[Links](#)

ELSEVIER
FULL-TEXT ARTICLE

Molecular cloning and binding properties of the human type II activin receptor.

Donaldson CJ, Mathews LS, Vale WW.

Clayton Foundation Laboratories for Peptide Biology, Salk Institute, La Jolla, CA 92037.

A full-length cDNA for the type II human activin receptor was cloned by hybridization from a human testis cDNA library. The sequence encodes a 513 amino acid protein that is 99% identical, at the amino acid level, with the mouse type II activin receptor. The type II human activin receptor consists of an extracellular domain that specifically binds activin A with a Kd of 360 pM, a single-membrane spanning domain, and an intracellular kinase domain with predicted serine/threonine specificity.

PMID: 1314589 [PubMed - indexed for MEDLINE]

Display Show Sort by Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Apr 24 2006 06:33:44